James J. Piscatelli, M.D. Jehad A. Miqdadi, M.D. Mohammad Fayyaz, M.D. Alok D. Gupta, M.D.

Mark Kuczmanski, M.S., PA-C Lindsay Harlock, PA-C Heather Glose, PA-C

# **GASTROENTEROLOGY**

Buffalo Medical Group, P.C.

## PREPARATION FOR FLEXIBLE SIGMOIDOSCOPY

You will need to purchase a 10oz bottle of Magnesium Citrate (over the counter) and Two Fleet enemas.

- <u>Do not take</u> any iron pills, iron supplements, and other supplements containing fish oil or flax seed oil 5 days before the exam.
- Please remember to continue taking asthma, anti-anxiety, or seizure medications!
- Please do not consume any corn, nuts or seeds for 5 days before the exam.

Please call our office to let us know if you are allergic to Versed or Fentanyl.

If you are a diabetic, please contact your Primary Care Physician. They will need to regulate the dosing of your insulin or glucose for the day of the procedure.

### THE DAY BEFORE THE PROCEDURE:

- 1. Have a light lunch at 12:00pm.
- 2. At 3 P.M. drink 10oz bottle of Magnesium Citrate
- 3. Have a clear liquid dinner that evening. (see below)
- 4. Do 1 Fleet enema before bed that evening.

#### THE DAY OF THE PROCEDURE:

- 1. Have a clear liquid breakfast ONLY.
- 2. Do 1 Fleet enema at 7:00am.

A **CLEAR LIQUID** diet means: no milk or milk products; no solid foods; no products containing RED or PURPLE dyes. Most cola soft drinks like Pepsi and Coke have red dyes in them.

You can have water, clear broth or bouillon, coffee/tea (without milk or non-dairy creamer), sherbet, strained fruit juices without pulp. Also the following if not RED or PURPLE: Gatorade, ice popsicles, carbonated and non-carbonated soft drinks, Kool-Aid, plain Jell-O (without fruit or toppings).

REPORT TO:			
TIME:			
DATE:			

If you are currently taking any of the medications listed on the next page, please contact your prescribing physician for clearance to stop these medications.

Mark Kuczmanski, M.S., PA-C Lindsay Harlock, PA-C Heather Glose, PA-C

# **GASTROENTEROLOGY**

Buffalo Medical Group, P.C.

If you are currently taking any of the medications listed below, please contact your prescribing physician for clearance to stop these medications.

# **Anticoagulation Therapy**

## **Antiplatelets:**

Plavix® (clopidogrel) Brilinta® (ticagrelor) Effient® (prasugrel) Aspirin

#### Other NSAIDs:

diclofenac, ibuprofen (Motrin®), indomethacin, ketoprofen delecoxib (Celebrex®), diflunisal, naproxen (Aleve®, Naprosyn®), sulindac meloxicam (Mobic®), nabumetone, piroxicam (Feldene®)

### **Anticoagulants:**

Coumadin® (warfarin) Eliquis® (apixaban) Pradaxa® (dabigatran) Xarelto® (rivaroxaban) Savaysa® (endoxaban)